



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Journal Pre-proofs

Clinical study


Prevalence, severity, outcomes, and risk factors of COVID-19 in multiple sclerosis: An observational study in the Middle East

Raed Alroughani, Jihad Inshasi, Jasem Al-Hashel, Jaber Alkhaboury, Abdullah Alsalti, Reem Al Suwaidi, Loqman H. Hassino, Samar Farouk Ahmed

PII: S0967-5868(22)00086-8
DOI: <https://doi.org/10.1016/j.jocn.2022.02.033>
Reference: YJOCN 9961

To appear in: *Journal of Clinical Neuroscience*

Received Date: 9 August 2021
Revised Date: 10 February 2022
Accepted Date: 22 February 2022



Please cite this article as: R. Alroughani, J. Inshasi, J. Al-Hashel, J. Alkhaboury, A. Alsalti, R. Al Suwaidi, L.H. Hassino, S. Farouk Ahmed, Prevalence, severity, outcomes, and risk factors of COVID-19 in multiple sclerosis: An observational study in the Middle East, *Journal of Clinical Neuroscience* (2022), doi: <https://doi.org/10.1016/j.jocn.2022.02.033>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Study Title

Prevalence, severity, outcomes, and risk factors of COVID-19 in multiple sclerosis: An observational study in the Middle East.

Raed Alroughani¹, Jihad Inshasi², Jasem Al-Hashel³, Jaber Alkhaboury⁴, Abdullah Alsalti⁵, Reem Al Suwaidi⁶, Loqman H. Hassino⁷, Samar Farouk Ahmed⁸

¹Division of Neurology,
Department of Medicine, Amiri Hospital,
Sharq, Kuwait
alroughani@gmail.com

²Neurology Department,
Rashid Hospital and Dubai Medical College,
Dubai Health Authority (DHA), Dubai, UAE
inshasi@emirates.net.ae

³Consultant Neurologist and Associate Professor of Neurology
Faculty of Medicine, Kuwait University, Kuwait
jasemkumsa@hotmail.com

⁴Neurology Unit, College of Medicine and Health Sciences,
Sultan Qaboos University and Sultan Qaboos University Hospital,
Muscat, Oman
jaberoman@hotmail.com

⁵Neurology Unit, College of Medicine and Health Sciences,
Sultan Qaboos University and Sultan Qaboos University Hospital,
Muscat, Oman
dr_alsalti@hotmail.com

⁶Neurology Department,
Rashid Hospital and Dubai Medical College,
Dubai Health Authority (DHA), Dubai, UAE
Dr_alsuwaidi@live.com

⁷Neurology Unit,
Department of Medicine Jahra Hospital,
Jahra City, Kuwait
lhassino@yahoo.com

⁸Senior Specialist Neurology, Ibn Sina Hospital, Kuwait,
Professor in Neurology, Minia University, Egypt
samerelshayb@hotmail.com

Corresponding Author:

Dr. Raed Alroughani
Division of Neurology,
Department of Medicine, Amiri Hospital,
Sharq, Kuwait
alroughani@gmail.com
Tel: +96599304949
Fax: +965 9779 6981

Declaration of Interest: None

Funding Sources

The development of this publication was financially supported by Merck Serono Middle East FZ Ltd., an affiliate of Merck KGaA, Darmstadt, Germany through an independent medical writing grant. The views and opinions expressed in this publication do not necessarily reflect those of the grantor.

Abstract

A cross-sectional hospital records-based study was conducted to evaluate the prevalence, severity, outcomes, and identify demographic and clinical risk factors of coronavirus disease (COVID-19) in patients with MS. The study was conducted at multiple clinics in Oman, Kuwait, and the United Arab Emirates (UAE) from March 2020 to February 2021. The association of patient demographics, MS disease characteristics, and use of disease-modifying therapies with outcomes of COVID-19 illness were evaluated using odds ratio. A total of 134 MS patients with COVID-19 (prevalence rate of 3.7%) having a median age of 35.5 years were analyzed in the study. A majority (126 [94.0%]) of patients had mild COVID-19 illness and 122 (91.0%) made a full recovery, while 1 (0.7%) patient died. The median EDSS score reported in the study was low (1.0). Univariate regression analysis showed high EDSS scores, progressive MS disease, and use of anti-CD20 therapy such as rituximab as risk factors for moderate to severe COVID-19 requiring hospitalization. Comorbidities were associated with a higher risk of non-recovery from COVID-19 in both univariate and multivariate analyses. Age, sex, smoking history, and duration of MS did not show a significant association with severity or adverse COVID-19 disease outcome. Identification of risk factors can aid in improving the treatment and monitoring of pwMS and COVID-19.

Keywords

COVID-19; disease-modifying therapy; multiple sclerosis; regression analyses, odds ratio; anti- CD20 therapy; comorbidity

Introduction

The emergence of the coronavirus disease 2019 (COVID-19) pandemic has raised substantial concerns regarding its impact on populations affected by auto-immune diseases. MS, an autoimmune neurodegenerative disorder of the central nervous system, is being studied for its correlation with COVID-19. [1, 2, 3]

MS patients are considered a high-risk population for COVID-19, due to the high prevalence of disability and widespread use of immunosuppressive disease-modifying therapy (DMT). [4] Among MS patients, those who belong to an older age group or have a more progressive form of the disease have been shown to develop complications in addition to respiratory tract symptoms during a COVID-19 infection. [5]

Indeed, MS patients are found to be at a greater risk of exacerbations and relapses due to this viral infection. [4, 6]

The exact relationship between COVID-19 and MS is still unclear. There are contradictory reports regarding the relation between COVID-19 severity/risk and immunosuppressive medications. The primary aim of our study was to evaluate the prevalence, severity, and outcomes of COVID-19 and to identify demographic and clinical risk factors in pwMS from three countries in the Middle East, namely Oman, Kuwait, and the United Arab Emirates.

Methodology

This was a retrospective, cross-sectional, observational study conducted in one MS clinic in Oman (Khaoula hospital, Muscat), three clinics in Kuwait (Ibn Sina hospital, Amiri hospital, Jahra hospital), and one clinic in the United Arab Emirates (Rashid hospital, Dubai). The study was conducted from March 2020 to February 2021. The data was collected from hospital records. MS was diagnosed according to the revised 2010 McDonald criteria. Before the application of these criteria, the diagnosis of MS was based on previously accepted diagnostic criteria. Demographic data including patient's age, gender, date of MS onset (date of first clinical event) and diagnosis, presentation at onset, disease duration (time in years since the first clinical event), type of MS, expanded disability status scale (EDSS) scores, and the use of disease-modifying therapies (DMTs) was extracted from the records. Data of MS patients who were diagnosed with COVID-19 were also extracted from the records. The data were confirmed by either contacting patients

by phone or when they visited the clinics for their routine scheduled visits or due to their concern about their DMTs. The patients were asked to complete a self-administered survey sent via email or filled at the clinic that retrospectively collected details of the course of their COVID-19 illness such as symptoms, investigations performed, and requirement of hospitalization. Information was also obtained from their medical records. The severity of COVID-19 was categorized as mild (not requiring hospitalization), moderate (hospitalized), and severe (requiring intensive care). MS patients older than 18 years of age with the following diagnoses were included: clinically isolated syndrome (CIS), relapsing-remitting MS (RRMS), secondary progressive MS (SPMS), or primary progressive MS (PPMS). Diagnosis of COVID-19 was done either through PCR, serology, or chest imaging. The endpoints for this study were the prevalence rate of COVID-19 in the studied MS cohort and measures of association of COVID-19 severity, hospitalization, and outcome with key demographic and clinical variables of MS.

Statistical Analysis

The prevalence of COVID-19 in MS patients was estimated by dividing the number of confirmed COVID-19 cases in the cohort by the total number of MS patients as per the hospital records. This was compared to the COVID-19 prevalence in the general population of that country/city. Descriptive statistics were used to compare demographics, disease characteristics, and the severity of MS. Univariate logistic regression models were performed on identified variables, namely, COVID-19 outcome (recovery or non-recovery), COVID-19 severity (mild or non-mild [moderate/severe]), and hospital admission (Yes/No) to assess their association with covariates such as age (< 45 or ≥ 45), gender, smoking, EDSS score (0-4.5 or 5-9.5), comorbidities, MS diagnosis (RRMS/CIS or progressive [SPMS/PPMS]), MS disease duration, (≤ 10 years or > 10 years), On/Off DMTs, duration of current DMT (≤ 5 years or > 5 years), and type of DMT. Pearson's chi-squared test, Fisher's exact test, Student's t-test, and the Mann–Whitney U test were used for comparisons as appropriate. Odds ratios (ORs) with 95% confidence intervals (CIs) were estimated with a logistic regression model. A multivariate logistic regression analysis was performed using the same variables used in the univariate analysis.

Results

As of February 26, 2021, a total of 3624 pwMS were reported in the records of previously specified clinics and hospitals of Kuwait (2334), Dubai, UAE (1050), and Oman (240). Of

these, 134 patients (Kuwait: 92 (3.9%), Dubai, UAE: 36 (3.4%), Oman: 6 (2.5%)) had tested positive for COVID-19 with an overall prevalence rate of 3.7% in the total analyzed MS population. Demographics and clinical characteristics of the cohort of 134 patients are presented in **Table 1**. Ninety-eight patients (73.1%) were female, with the median age of the cohort being 35.5 years, and a mean (SD) disease duration of 8.83 (6.0) years. Non-smokers accounted for 88.8% of the total cohort with 91% reporting no morbid obesity. Relapsing-remitting MS accounted for 86.6% of all types of MS patients analyzed in the cohort. The majority (90.3%) had no comorbidities. Hypertension/cardiovascular disease (5 [3.7%]), malignancy in remission (2 [1.5%]), renal disease (1 [0.7%]), and liver disease (1 [0.7%]) were few co-morbidities reported. The median EDSS score (1.0) was in the range of 0-4.5 for 127 (94.8%) patients while 7 (5.2%) patients reported a score in the range of 5-9.5 [Table 1].

A total of one twenty-eight (95.5%) patients were diagnosed by polymerase chain reaction, four (3.0%) patients had a positive serology, and one (0.7%) patient had chest imaging showing typical lesions for COVID-19, while the diagnostic methodology was unknown for one patient. Overall, 127 (94.8%) patients were on DMTs; the four most common DMTs being used were fingolimod (26.9%), ocrelizumab (15.7%), and rituximab and natalizumab (10.4% each), with 5.2% reporting no MS treatment [Table1]. The majority (94.0%) of patients reported mild COVID-19, 3.7% reported moderate severity of infection, while 1.5% reported severe COVID-19, and one (0.7%) patient was in a life-threatening condition. Patients presented with typical symptoms for COVID-19. Notably, fever (temperature more than 38 °C), fatigue, dry cough, and sore throat were more frequent in patients, along with anosmia. Eight (6.0%) patients were hospitalized; of these, 3 (2.2%) were in the intensive care unit and 2 (1.5%) required ventilator support. In evaluating COVID-19 outcome in terms of recovery and non-recovery, we found that 122 (91.0%) patients recovered, and one (0.7%) patient died. At the time of writing this manuscript, 9 (6.7%) patients had an ongoing COVID-19 infection, and the outcome was unknown in 2 (1.5%) patients.

To estimate the risk factors associated with COVID-19 severity, hospitalization, and outcomes, we calculated the odds ratios (ORs) from univariate and multivariate logistic regression models. Since all moderately ill patients were also hospitalized, the statistical results of the association of various covariates were similar for both, hospitalization, and COVID-19 severity. Univariate analysis showed a statistically significant higher risk of

moderate/severe COVID-19 severity and hospitalization in patients with higher EDSS scores (OR: 8.06; 95% CI 1.290 – 50.453, $p = 0.026$) and in patients with progressive disease (OR 8.85; 95% CI 1.786 – 43.855, $p = 0.008$) [Table 2]. The association between age and COVID-19 severity was found to be non-significant ($p = 0.082$) with a 3.8 times higher risk of moderate/severe disease and hospitalization in patients aged 45 and above.

The presence of at least one comorbidity was significantly associated with a 5.3 times higher risk of non-recovery from COVID-19 (95% CI 1.174 – 24.225, $p = 0.03$) [Table 3]. Univariate analysis of DMT use showed a statistically significant higher risk of developing moderate/severe COVID-19 and consequent hospitalization in patients on rituximab (6.2 times higher risk) (95% CI 1.319 – 29.830, $p = 0.021$). However, patients on other DMTs including ocrelizumab did not show an association with COVID-19 severity. The category of anti-CD20 therapy (rituximab and ocrelizumab combined data) showed significant association with COVID-19 severity (5.3 times higher risk) (95% CI 1.203 – 23.639, $p = 0.028$) [Table 4]. There was no association between DMT use and the outcome of COVID-19 ($p > 0.05$).

Multivariate regression analysis did not show a significant association of any of the covariates tested with the severity of COVID-19 or hospitalization [Table 5]. However, similar to univariate analysis, a significant association of non-recovery was found with the presence of comorbidities. There was a significant 1.5 times higher risk of non-recovery in patients with any comorbid condition, eliminating the effect of all other covariates (95% CI 1.073 – 2.134, $p = 0.018$) [Table 6].

Discussion

This hospital records-based observational study analyzed cohorts of MS patients from three clinics in Kuwait and one clinic each in UAE and Oman for eleven months. The prevalence rate of COVID-19 (3.7%) in the analyzed MS cohort is marginally lower than the overall prevalence of COVID-19 in the general population of Kuwait (5.3%), Dubai (4.75%), and Oman (4.07%). [7] A few similar studies investigating the effects of COVID-19 in MS patients have been reported in literature. A study early on in the pandemic carried out in Italy reported similar prevalence of COVID-19 among MS patients as in the general population. [8] Zabalza et al. reported on the incidence of COVID-19 in MS patients followed at a center in Barcelona, Spain. The incidence of confirmed COVID-19 among MS patients (6.3%) was found to be similar to that in the general population of the Catalan

region of Spain (6.1%). [1] Similarly, Sepulveda et al conducted a cross-sectional study to investigate the incidence of COVID-19 in a cohort of MS patients at a single center in Barcelona, Spain. The study found that the cumulative incidence of confirmed COVID-19 cases (1.27%) was similar to that of the general population (1.32%). [3] The results of these studies corroborate our findings of similar or only slightly lower prevalence of COVID-19 in MS patients compared to that in the general population of that region or country.

The hospitalization rate (6.0%), the requirement of intensive care (2.2%), and the mortality rate (0.7%) in our cohort were low. It could be possible that the younger age (median age, 35.5 years) of the patients with no comorbidities being reported in a majority of them (90.3%) and their low physical disability status (median EDSS score 1.0) may have contributed to mitigating their COVID-19 course. The lower rates of hospitalization and deaths in our study are also similar to those reported in other studies that have a low percentage of confirmed cases of COVID-19 in MS patients. For example, a UK community-based study with 15.6% of confirmed cases of COVID-19 in MS patients reported 1.3% of hospitalizations and no deaths [9]; a European and US study with 17% of confirmed cases of COVID-19 in MS patients described 5.6% of hospitalizations and no deaths [10], and an Italian program with 24.5% of confirmed cases reported 4% of patients with mild disease and 2.1% of mortality [11]. In contrast, higher rates of hospitalization (21.0%–25.6%) were found in studies with a higher proportion of confirmed cases. [5, 12, 13]

Clinical characteristics of COVID-19 in MS patients in our study do not differ greatly from those in the general population and are similar to those seen in other published articles. [14, 15, 16] In our cohort, fever, fatigue, dry cough, and sore throat were the most common symptoms.

In univariate analyses, higher EDSS scores (≥ 5) and progressive disease were identified as risk factors for increased COVID-19 severity and consequent hospitalization, while risk factors for an unfavorable outcome in terms of recovery and non-recovery were the presence of comorbid conditions. A similar multicenter, retrospective, observational cohort study was conducted by Louapre et al. to evaluate risk factors and outcomes of COVID-19 in MS patients in France. The study used the Covisep registry and identified greater age, male sex, comorbidities, progressive disease, and higher EDSS as risk factors for increased COVID-19 severity in their univariate analyses. [5]

In the general population, the risk for severe COVID-19 illness increases with age, and older adults are at the highest risk. [16] In our study, we found age to be a risk factor with 3.8 times higher risk of moderate/severe COVID-19 in MS patients who are above the age of 45 years, but this association was not statistically significant. This may be attributed to a lower number of patients (20) in the older age group (≥ 45) in our cohort. In the study conducted by Parotta et al., critically ill COVID-19 infected MS patients were older (mean age of 57.7 years). [12]

It is established that respiratory dysfunction is common in pwMS an EDSS of 7 or more and is linked to impairment of expiratory muscles. [17] In our univariate analyses, EDSS was an independent risk factor for a severe form of COVID-19. The cohort we analyzed had a low level of neurological disability as assessed by EDSS scores, with 94.8% reporting scores between 0 – 4.5; however, higher disability scores [5 – 9.5] had a significant association with increasing severity of COVID-19 infection. The association between COVID-19 severity and EDSS has been previously reported only by Louapre et al.; COVID-19 patients with severity scores of 3 or more (moderate/severe) on the COVID-19 ordinal scale had higher EDSS (median EDSS, 6.0). [5] The identification of higher EDSS as a risk factor for severe COVID-19 and increased risk of hospitalization should lead to the adoption of sufficient and sustainable precautionary measures for limiting COVID-19 infection in these patients.

Certain comorbid conditions in COVID-19 are known to increase the risk for severe illness. pwMS frequently have several comorbidities that have been related to a worse COVID-19 prognosis. [5] Univariate and multivariate analysis in our study also showed the presence of co-morbidities as an independent risk factor for COVID-19 outcome with a higher chance of non-recovery. Due to the insufficient number of patients, we were unable to establish the association of specific comorbidities with COVID-19 severity or outcome.

Relapsing-remitting MS is typically known to be the most common type of MS and most patients are diagnosed initially with this type. The disease takes a more progressive course after several decades. The majority of patients in our cohort had RRMS (86.6%). Considering the lower age of the participants, this is understandable. Progressive disease was associated with increased COVID-19 severity in our cohort. This was similar to the results of studies conducted by Louapre et al. and Parotta et al. that found progressive MS to be a risk factor for developing more severe COVID-19. [5, 12]

Disease-modifying therapies are known to modify the risk of COVID-19 infection based on their mechanisms of action. In our study, we found a statistically significant higher risk of developing moderate/severe COVID-19 and consequent hospitalization in patients on rituximab (6.3 times higher risk) and anti-CD20 therapy (5.3 times higher risk). However, a few studies have found that the type/class of DMT is not a predictor for increased severity of COVID-19 or of poor outcome. [3, 5, 8, 12, 18, 19] A large electronic health record-based observational study conducted by Reder et al. in 2021 found that pwMS taking interferons and glatiramer acetate had the lowest risk of COVID-19 whereas patients receiving anti-CD20 therapies had the highest risk. [20] Another study by Fragoso et al. carried out using data from five Latin American countries in patients with MS and confirmed diagnosis of COVID-19 reported the use of anti-CD20 therapies as the only risk factor associated with hospitalization and death. [21] Other studies including one with a large multinational sample of MS patients have also implicated anti-CD20 therapy (either rituximab or ocrelizumab) in increasing the risk of severe COVID-19 and hospitalization/ICU admission. [22, 23, 24] The results of these studies corroborate our results and can be explained by the fact that anti-CD20 therapies impact the humoral immune response and decrease the levels of IgG and IgM over time. [25] However, any association with DMTs should be inferred with caution as results may be attributed to inadequate sample size and may vary if a larger cohort with the use of any of the DMTs is analyzed. A study by Zabalza et al. [1] found longer MS disease duration to be a risk factor for COVID-19; however, this association was not seen in our study.

Our study has several notable limitations that include the lack of a structured pre-defined data collection format; it also involved voluntary reporting by health care professionals, which may have led to biased reporting towards more severe cases. Considering the retrospective nature of the study there may also be some amount of recall bias. The small sample size does not let an appropriate assessment of the DMT-related COVID-19 risk of infection. The study did not use the WHO-recommended COVID-19 eight-category ordinal scale for describing disease severity. The lack of a non-infected cohort makes it difficult to make any assumptions and comparisons between MS characteristics in COVID-19 patients and among those in the general MS population. An adequate number of patients were not included in the analysis in Oman due to the lack of a comprehensive follow-up protocol and the prevailing COVID scenario. The strength of the study is that it is one of the few studies on MS patients with COVID-19 from the Middle East and can provide

valuable insights on risk factors leading to more severe disease and consequent adverse outcomes in this patient population.

Conclusions

The prevalence of COVID-19 in the MS patient cohort from the Middle East is marginally lower than that in the general population. COVID-19 severity and outcome in this cohort are largely dependent on a few key indicators such as the presence of co-morbidities, higher scores of disabilities, presence of progressive forms of MS, and the use of anti-CD 20 DMTs. The prognosis of COVID-19 is generally favorable in a majority of COVID-19 pwMS.

Acknowledgments

Dr. Rupali Bahri from Medcytes, Dubai provided medical writing services for the development of this manuscript.

References

1. Zabalza A, Cárdenas-Robledo S, Tagliani P, Arrambide G, Otero-Romero S, Carbonell-Mirabent P, et al. COVID-19 in multiple sclerosis patients: susceptibility, severity risk factors and serological response. *Eur J Neurol*. 2020 [ahead of print]. doi:10.1111/ene.14690.
2. Salter A, Fox RJ, Newsome SD, Halper J, Li DKB, Kanellis P, et al. Outcomes and risk factors associated with SARS-CoV-2 infection in a North American registry of patients with multiple sclerosis. *JAMA Neurol*. 2021;78(6):699–708. doi:10.1001/jamaneurol.2021.0688.
3. Sepúlveda M, Llufríu S, Martínez-Hernández E, Català M, Artola M, Hernando A, et al. Incidence and impact of COVID-19 in MS: A survey from a Barcelona MS unit. *Neurol Neuroimmunol Neuroinflamm*. 2021;8(2):e954. doi:10.1212/NXI.0000000000000954.
4. Choudhury F, Bulka H, Rathnam AS, Said OM, Lin J, Lorigan H, et al. COVID-19 in multiple sclerosis patients and risk factors for severe infection. *J Neurol Sci*. 2020;418:117147. doi: 10.1016/j.jns.2020.117147.
5. Louapre C, Collongues N, Stankoff B, Giannesini C, Papeix C, Bensa C, et al. Clinical characteristics, and outcomes in patients with coronavirus disease 2019 and multiple sclerosis. *JAMA Neurol*. 2020;77(9):1079-1088. doi:10.1001/jamaneurol.2020.2581.

6. Sadeghmousavi S, Rezaei N. COVID-19 and multiple sclerosis: predisposition and precautions in treatment. *SN Compr Clin Med*. 2020;1-6. doi:10.1007/s42399-020-00504-9.
7. Worldometer real-time world statistics: Corona update live <https://www.worldometers.info/coronavirus/>; 2021[accessed 03 March 2021].
8. Mantero V, Abate L, Balgera R, Basilico P, Salmaggi A, Cordano C. Assessing the susceptibility to acute respiratory illness COVID-19-related in a cohort of multiple sclerosis patients. *Mult Scler Relat Disord*. 2020;46:102453. doi: 10.1016/j.msard.2020.102453.
9. Evangelou N, Garjani A, dasNair R, Hunter R, Tuite-Dalton KA, Craig EM, et al. Self-diagnosed COVID-19 in people with multiple sclerosis: a community-based cohort of the UK MS register. *J Neurol Neurosurg Psychiatry*. 2020;92(1):107–109. doi: 10.1136/jnnp-2020-324449.
10. Moss BP, Mahajan KR, Bermel RA, Hellisz K, Hua LH, Hudec T, et al. Multiple sclerosis management during the COVID-19 pandemic. *Mult Scler*. 2020 26(10):1163-1171. doi: 10.1177/1352458520948231.
11. Crescenzo F, Marastoni D, Bovo C, Calabrese M. Frequency, and severity of COVID-19 in multiple sclerosis: A short single-site report from northern Italy. *Mult Scler Relat Disord*. 2020 Sep;44:102372. doi: 10.1016/j.msard.2020.102372.
12. Parrotta E, Kister I, Charvet L, Sammarco C, Saha V, Charlson RE, et al. COVID-19 outcomes in MS: Observational study of early experience from NYU Multiple Sclerosis Comprehensive Care Center. *Neurol Neuroimmunol Neuroinflamm*. 2020;7(5):e835. doi:10.1212/NXI.0000000000000835.
13. Loonstra FC, Hoitsma E, van Kempen ZLE, Killestein J, Mostert JP. COVID-19 in multiple sclerosis: The Dutch experience. *Mult Scler*. 2020;26(10):1256-1260. doi:10.1177/1352458520942198.
14. Safavi F, Nourbakhsh B, Azimi AR. B-cell depleting therapies may affect susceptibility to acute respiratory illness among patients with multiple sclerosis during the early COVID-19 epidemic in Iran. *Mult Scler Relat Disord*. 2020;43:102195. doi:10.1016/j.msard.2020.102195.
15. Sormani MP on behalf of the Italian Study Group on COVID-19 infection in multiple sclerosis. An Italian programme for COVID-19 infection in multiple sclerosis. *Lancet Neurol* 2020;19(6):481-482. doi:10.1016/S1474-4422(20)30147-2.

16. Centers for Disease Control: Coronavirus guidelines and precautions <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/older-adults.html>; 2021 [accessed 03 March 2021].
17. Levy J, Bensmail D, Brotier-Chomienne A, Butel S, Joussain C, Hugeron C, et al. Respiratory impairment in multiple sclerosis: A study of respiratory function in wheelchair-bound patients. *Eur J Neurol.* 2017;24(3):497-502. doi: 10.1111/ene.13231.
18. Sen S, Karabudak R, Schiavetti I, Demir S, Ozakbas S, Tutuncu M, et al. "The outcome of a national MS-Covid-19 study: What the Turkish MS cohort reveals?" *Mult Scler Relat Disord.* 2021 Jul;52:102968. doi:10.1016/j.msard.2021.102968.
19. Zakaria M, Ponzano M, Schiavetti I, Carmisciano L, Nada M, Abdel Nasser M, et al. The MuSC-19 study: The Egyptian cohort. *Mult Scler Relat Disord.* 2021;56:103324. doi: 10.1016/j.msard.2021.103324.
20. Reder AT, Centonze D, Naylor ML, Nagpal A, Rajbhandari R, Altincatal A, et al. COVID-19 in patients with multiple sclerosis: Associations with disease-modifying therapies. *CNS Drugs.* 2021;35(3):317-330. doi:10.1007/s40263-021-00804-1.
21. Fragoso YD, Schiavetti I, Carmisciano L, Ponzano M, Steinberg J, Treviño-Frenk I, et al. Coronavirus disease 2019 in Latin American patients with multiple sclerosis. *Mult Scler Relat Disord.* 2021 Oct;55:103173. doi: 10.1016/j.msard.2021.103173.
22. Sormani MP, De Rossi N, Schiavetti I, Carmisciano L, Cordioli C, Moiola L, et al. Disease modifying therapies and coronavirus disease 2019 severity in multiple sclerosis. *Ann Neurol.* 2021;89:780-789. doi: 10.1002/ana.26028.
23. Simpson-Yap S, De Brouwer E, Kalincik T, Rijke N, Hillert JA, Walton C, et al. Associations of disease-modifying therapies with COVID-19 severity in multiple sclerosis. *Neurology.* 2021;97:e1870-e1885. doi:10.1212/WNL.00000000000012753.
24. Landtblom A-M, Berntsson SG, Bostrom I, Iacobaeus E. Multiple sclerosis and COVID-19: The Swedish experience. *Acta Neurol Scand.* 2021;00:1–7. doi: 10.1111/ane.13453.
25. Zheng C, Kar I, Chen CK, Sau C, Woodson S, Serra A, et al. Multiple sclerosis disease-modifying therapy and the COVID-19 pandemic: Implications on the risk

of infection and future vaccination. *CNS Drugs*. 2020;34:879–896.
doi:10.1007/s40263-020-00756-y.

Table 1: The distribution of demographic and clinical variables for COVID-19 MS patients

Demographic and Clinical Characteristics		N = 134
Age	Median	35.5
	Mean (SD)	36.1 (9.7)
	< 45	114
	≥ 45	20
Sex, n (%)	Male	36 (26.9)
	Female	98 (73.1)
Smoking, n (%)	Yes	15 (11.2) ^a
Obesity, n (%)	Yes	12 (9.0) ^a
Pregnancy, n (%)	Yes	2 (1.5) ^a
	No	121 (90.3)
Comorbidities, n (%)	CVS/HTN	5 (3.7)
	Malignancy in remission	2 (1.5)
	Renal function	1 (0.7)
	Liver disease	1 (0.7)
	Others	1 (0.7)
Diagnosis, n (%)	CIS	7 (5.2)
	RRMS	116 (86.6)
	SPMS	7 (5.2)
	PPMS	4 (3.0)
Disease Duration	Mean (SD)	8.83 (6.0)
	≤ 10 years	91 (67.9)
	> 10 years	43 (32.1)
On/Off DMT, n (%)	Off DMT	7 (5.2)
	On DMT	127 (94.8)
Current DMT, n (%)	No	7 (5.2)
	IFNB	12 (9.0)
	GA	1 (0.7)
	Teriflunomide	7 (5.2)

	DMF	11 (8.2)
	Fingolimod	36 (26.9)
	Natalizumab	14 (10.4)
	Alemtuzumab	4 (3.0)
	Ocrelizumab	21 (15.7)
	Rituximab	14 (10.4)
	Cladribine tablets	4 (3.0)
	Others	3 (2.2)
Duration of Current DMT, n (%)	≤ 5 years	109 (81.3)
	> 5 years	23 (17.2)
Diagnosis of COVID-19, n (%)	PCR	128 (95.5)
	Serology	4 (3.0)
	Typical chest imaging	1 (0.7)
COVID-19 Symptoms, n (%)	No	12 (9.0)
	Yes	122 (91.0)
Symptoms, n (%)	Fever	105 (78.4)
	Fatigue	86 (64.2)
	Dry cough	74 (55.2)
	Sore throat	70 (52.2)
	Anosmia	47 (35.1)
	Pain	22 (16.4)
	Pneumonia	6 (4.5)
	Diarrhea	8 (6.0)
Severity, n (%) ^b	Mild	126 (94.0)
	Moderate	5 (3.7)
	Severe	2 (1.5)
	Life threatening	1 (0.7)
Hospital Admission, n (%)	Yes	8 (6.0)
ICU Admission, n (%)	Yes	3 (2.2) ^a
Ventilator Required, n (%)	Yes	2 (1.5)
ECMO required, n (%)	Yes	1 (0.7)
COVID-19 Outcomes, n (%)	Unknown	2 (1.5)
	Ongoing	9 (6.7)
	Recovery	122 (91.0)
	Death	1 (0.7)
EDSS Score, n (%)	0-4.5	127 (94.8)
	5-9.5	7 (5.2)
EDSS Score, median		1.0
Country, n (%)	Kuwait	92 (68.7)
	Dubai	36 (26.9)
	Oman	6 (4.5)

^a Contains some missing values.

^b Severity definitions: mild- those not requiring hospitalization, moderate- those requiring hospitalization, severe- those requiring admission to intensive care unit, life-threatening- those whose COVID-infection was life-threatening.

CIS, clinically isolated syndrome; COVID-19, coronavirus disease 2019; CVS, cardiovascular system; DMF, dimethyl fumarate; DMT, disease modifying therapy; ECMO, extracorporeal membrane oxygenation; EDSS, Expanded Disability Severity Scale; GA, glatiramer acetate; HTN, hypertension; ICU, intensive care unit; IFNB, interferon beta; MS, multiple sclerosis; PCR, polymerase chain reaction; PPMS, primary progressive multiple sclerosis; RRMS, relapsing-remitting multiple sclerosis; SPMS, secondary progressive multiple sclerosis.

Table 2: Summary of univariate logistic regression models for the COVID-19 MS patients in relation to COVID-19 severity

Variables	Categories	Odds Ratio	95% CI		P-Value ^a
			Lower Bound	Upper Bound	
Age Group	≥ 45	3.847	0.842	17.587	0.082
	< 45				
Gender	Female	0.591	0.134	2.612	0.488
	Male				
Smoker	Yes	1.133	0.130	9.898	0.910
	No				
EDSS	High (5-9.5)	8.067	1.290	50.453	0.026
	Low (0-4.5)				
Comorbidities	Any co-morbid condition	2.130	0.231	19.668	0.505
	No comorbidity				
MS Diagnosis	Progressive disease (PPMS+SPMS)	8.850	1.786	43.855	0.008
	Non-progressive disease (RRMS+CIS)				
Current DMT Duration	> 5 years	0.662	0.078	5.660	0.707
	≤ 5 years				
Disease Duration	> 10 years	1.290	0.294	5.65	0.736
	≤ 10 years				

^ap value is significant at <0.05

CI, confidence interval; CIS, clinically isolated syndrome; COVID-19, coronavirus disease 2019; DMT, disease modifying therapy; EDSS, Expanded Disability Severity Scale; MS, multiple sclerosis; PPMS, primary progressive multiple sclerosis; RRMS, relapsing-remitting multiple sclerosis; SPMS, secondary progressive multiple sclerosis.

Table 3: Summary of univariate logistic regression models for the COVID-19 MS patients in relation to COVID-19 outcome

Variables	Categories	Odds Ratio	95% CI		P-Value ^a
			Lower Bound	Upper Bound	
Age Group	≥ 45	1.156	0.234	5.715	0.859
	< 45				
Gender	Female	0.711	0.200	2.523	0.598
	Male				
Smoker	Yes	0.695	0.083	5.797	0.737
	No				
EDSS	High (5-9.5)	1.758	0.194	15.949	0.616
	Low (0-4.5)				
Comorbidities	Any co-morbid condition	5.333	1.174	24.225	0.030
	No comorbidity				
MS Diagnosis	Progressive disease (PPMS+SPMS)	2.511	0.476	13.247	0.278
	Non-progressive disease (RRMS+CIS)				
On/Off DMT	On DMT	0.569	0.063	5.163	0.616
	Off DMT				
Current DMT Duration	> 5 years	2.658	0.727	9.718	0.139
	≤ 5 years				
Disease Duration	> 10 years	2.297	0.695	7.594	0.173
	≤ 10 years				

^ap value is significant at <0.05.

CI, confidence interval; CIS, clinically isolated syndrome; COVID-19, coronavirus disease 2019; DMT, disease modifying therapy; EDSS, Expanded Disability Severity Scale; MS, multiple sclerosis; PPMS, primary progressive multiple sclerosis; RRMS, relapsing-remitting multiple sclerosis; SPMS, secondary progressive multiple sclerosis.

Table 4: Summary of univariate logistic regression models for the current DMT use among the COVID-19 MS patients in relation to COVID-19 severity

Variables ^b	Odds Ratio	95% CI		P-Value ^a
		Lower Bound	Upper Bound	
Fingolimod	0.371	0.044	3.129	0.362
Natalizumab	1.242	0.141	10.903	0.845
Ocrelizumab	1.877	0.352	10.002	0.461
Rituximab	6.273	1.319	29.830	0.021
Anti-CD20 (rituximab + ocrelizumab)	5.333	1.203	23.639	0.028

^ap value is significant at <0.05.

^bFour DMDs most commonly used by patients in the study have been listed over here. DMDs not listed in this table have a low number of patients and give skewed results.

CD20, cluster of differentiate 20; COVID-19, coronavirus disease 2019; DMT, disease modifying therapy; MS, multiple sclerosis.

Table 5: Summary of multivariate logistic regression models for the COVID-19 MS patients in relation to severity

Variables	Categories	Odds Ratio	95% CI		P-Value ^a
			Lower Bound	Upper Bound	
Age Group	≥ 45	1.797	0.217	14.899	0.587
	< 45				
Gender	Female	0.520	0.075	3.629	0.509
	Male				
Smoker	Yes	0.728	0.052	10.297	0.815
	No				
EDSS	High (5-9.5)	7.828	0.182	336.513	0.284
	Low (0-4.5)				
Comorbidities	Any co-morbid condition	1.218	0.751	1.977	0.424
	No comorbidity				
MS Diagnosis	Progressive disease (PPMS+SPMS)	1.007	0.033	30.687	0.997
	Non-progressive disease (RRMS+CIS)				
Current DMT Duration	> 5 years	1.100	0.099	12.246	0.938
	≤ 5 years				
Disease Duration	> 10 years	0.962	0.131	7.049	0.970
	≤ 10 years				

^ap value is significant at <0.05.

CI, confidence interval; CIS, clinically isolated syndrome; COVID-19, coronavirus disease 2019; DMT, disease modifying therapy; EDSS, Expanded Disability Severity Scale; MS, multiple sclerosis; PPMS, primary progressive multiple sclerosis; RRMS, relapsing-remitting multiple sclerosis; SPMS, secondary progressive multiple sclerosis.

Table 6: Summary of multivariate logistic regression models for the COVID-19 MS patients in relation to COVID-19 outcome

Variables	Categories	Odds Ratio	95% CI		P-Value ^a
			Lower Bound	Upper Bound	
Age Group	≥ 45	0.699	0.113	4.308	0.699
	< 45				
Gender	Female	0.564	0.125	2.536	0.455
	Male				
Smoker	No	0.408	0.038	4.382	0.46
	Yes				
EDSS	High (5-9.5)	1.334	0.038	46.821	0.874
	Low (0-4.5)				
Comorbidities	Any co-morbid condition	1.513	1.073	2.134	0.018
	No comorbidity				
MS Diagnosis	Progressive disease (PPMS+SPMS)	2.632	0.155	44.709	0.503
	Non-progressive disease (RRMS+CIS)				
Current DMT Duration	> 5 years	3.295	0.689	15.751	0.135
	≤ 5 years				
Disease Duration	> 10 years	1.156	0.246	5.43	0.855
	≤ 10 years				

^ap value is significant at <0.05.

CI, confidence interval; CIS, clinically isolated syndrome; COVID-19, coronavirus disease 2019; DMT, disease modifying therapy; EDSS, Expanded Disability Severity Scale; MS, multiple sclerosis; PPMS, primary progressive multiple sclerosis; RRMS, relapsing-remitting multiple sclerosis; SPMS, secondary progressive multiple sclerosis.

Highlights:

- Prevalence of COVID-19 in the studied MS cohort was marginally lower to that in the general population.
- The COVID -19 disease course was mild and outcomes were mostly favorable.
- Most patients did not require hospitalization.
- High EDSS score, progressive MS disease, and anti CD20 therapy use were risk factors for moderate to severe disease requiring hospitalization.
- Comorbidities were associated with a higher risk of non-recovery from COVID-19.